

| Espécies | P 1 (1) | P 1 (2) | P 2 (1) | P 2 (2) | P 3 (1) | P 3 (2) | P 4 (1) | P 4 (2) | P 5 (1) | P 5 (2) | Total | AR |
|-----------------------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|-------|-------|
| <i>Rhea americana</i> | 0 | 0 | 2 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 7 | 0,15% |
| <i>Nothura maculosa</i> | 4 | 0 | 1 | 3 | 2 | 2 | 1 | 0 | 0 | 0 | 13 | 0,28% |
| <i>Rhynchotus rufescens</i> | 1 | 1 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 4 | 0,09% |
| <i>Chauna torquata</i> | 2 | 0 | 2 | 2 | 2 | 2 | 2 | 0 | 0 | 2 | 14 | 0,31% |
| <i>Dendrocygna viduata</i> | 0 | 0 | 0 | 0 | 5 | 0 | 48 | 0 | 0 | 0 | 53 | 1,16% |
| <i>Cairina moschata</i> | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Amazonetta brasiliensis</i> | 6 | 2 | 26 | 17 | 5 | 7 | 22 | 16 | 0 | 2 | 103 | 2,25% |
| <i>Anas georgica</i> | 0 | 0 | 4 | 3 | 0 | 0 | 2 | 2 | 0 | 0 | 11 | 0,24% |
| <i>Anas flavirostris</i> | 3 | 0 | 9 | 0 | 0 | 0 | 22 | 35 | 0 | 0 | 69 | 1,51% |
| <i>Penelope obscura</i> | 3 | 0 | 4 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 9 | 0,20% |
| <i>Podilymbus podiceps</i> | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 2 | 0,04% |
| <i>Ciconia maguari</i> | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 2 | 0 | 0 | 3 | 0,07% |
| <i>Mycteria americana</i> | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 4 | 0 | 6 | 0,13% |
| <i>Phalacrocorax brasilianus</i> | 0 | 0 | 1 | 2 | 0 | 0 | 0 | 1 | 0 | 0 | 4 | 0,09% |
| <i>Ardea alba</i> | 1 | 0 | 2 | 1 | 0 | 1 | 0 | 1 | 4 | 0 | 10 | 0,22% |
| <i>Ardea cocoi</i> | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 2 | 0,04% |
| <i>Bubulcus ibis</i> | 0 | 0 | 2 | 1 | 0 | 1 | 29 | 0 | 143 | 0 | 176 | 3,85% |
| <i>Tigrisoma lineatum</i> | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Syrigma sibilatrix</i> | 2 | 3 | 4 | 0 | 4 | 2 | 1 | 0 | 0 | 0 | 16 | 0,35% |
| <i>Egretta caerulea</i> | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Theristicus caudatus</i> | 6 | 1 | 5 | 2 | 4 | 5 | 0 | 0 | 0 | 0 | 23 | 0,50% |
| <i>Plegadis chihi</i> | 0 | 0 | 33 | 0 | 6 | 0 | 60 | 0 | 0 | 13 | 112 | 2,45% |
| <i>Phimosus infuscatus</i> | 0 | 0 | 10 | 5 | 5 | 0 | 4 | 0 | 43 | 0 | 67 | 1,46% |
| <i>Theristicus caerulescens</i> | 0 | 0 | 2 | 2 | 0 | 1 | 3 | 2 | 0 | 0 | 10 | 0,22% |
| <i>Platalea ajaja</i> | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 10 | 0 | 0 | 11 | 0,24% |
| <i>Cathartes aura</i> | 1 | 9 | 1 | 3 | 2 | 1 | 4 | 13 | 0 | 0 | 34 | 0,74% |
| <i>Cathartes burrovianus</i> | 0 | 0 | 3 | 0 | 4 | 0 | 0 | 0 | 0 | 0 | 7 | 0,15% |
| <i>Coragyps atratus</i> | 1 | 3 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 5 | 0,11% |
| <i>Elanus leucurus</i> | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Heterospizias meridionalis</i> | 1 | 0 | 1 | 0 | 2 | 1 | 1 | 0 | 0 | 0 | 6 | 0,13% |

| | | | | | | | | | | | | |
|------------------------------|----|----|----|----|-----|-----|-----|-----|----|----|-----|--------|
| <i>Rostramus sociabilis</i> | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Rupornis magnirostris</i> | 1 | 0 | 2 | 1 | 2 | 1 | 0 | 0 | 0 | 0 | 7 | 0,15% |
| <i>Circus buffoni</i> | 0 | 0 | 2 | 1 | 4 | 2 | 0 | 0 | 0 | 0 | 9 | 0,20% |
| <i>Accipiter striatus</i> | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 0 | 0 | 0 | 4 | 0,09% |
| <i>Urubitinga urubitinga</i> | 0 | 0 | 0 | 0 | 1 | 0 | 7 | 0 | 1 | 0 | 9 | 0,20% |
| <i>Milvago chimango</i> | 2 | 0 | 1 | 1 | 5 | 2 | 17 | 0 | 0 | 0 | 28 | 0,61% |
| <i>Milvago chimachima</i> | 0 | 1 | 2 | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 6 | 0,13% |
| <i>Falco sparverius</i> | 1 | 1 | 2 | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 8 | 0,17% |
| <i>Caracara plancus</i> | 5 | 2 | 7 | 3 | 1 | 2 | 1 | 0 | 0 | 0 | 21 | 0,46% |
| <i>Aramus guarauna</i> | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Aramides ypecaha</i> | 2 | 0 | 4 | 3 | 4 | 2 | 0 | 1 | 0 | 0 | 16 | 0,35% |
| <i>Gallinula galeata</i> | 0 | 0 | 0 | 0 | 4 | 0 | 0 | 0 | 0 | 0 | 4 | 0,09% |
| <i>Gallinula melanops</i> | 0 | 0 | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 3 | 0,07% |
| <i>Cariama cristata</i> | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Vanellus chilensis</i> | 8 | 4 | 21 | 13 | 17 | 5 | 7 | 2 | 1 | 3 | 81 | 1,77% |
| <i>Himantopus melanurus</i> | 0 | 0 | 4 | 0 | 0 | 3 | 4 | 3 | 0 | 0 | 14 | 0,31% |
| <i>Gallinago paraguaiiae</i> | 0 | 0 | 1 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 0,09% |
| <i>Calidris sp</i> | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 0 | 0 | 3 | 0,07% |
| <i>Calidris melanotos</i> | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 0 | 0 | 0 | 3 | 0,07% |
| <i>Jacana jacana</i> | 0 | 0 | 2 | 0 | 4 | 0 | 0 | 2 | 0 | 0 | 8 | 0,17% |
| <i>Columba livia</i> | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 0,07% |
| <i>Columbina talpacoti</i> | 5 | 3 | 4 | 3 | 7 | 4 | 4 | 0 | 0 | 0 | 30 | 0,66% |
| <i>Columbina picui</i> | 7 | 3 | 7 | 5 | 3 | 5 | 8 | 15 | 5 | 6 | 64 | 1,40% |
| <i>Leptotila verreauxi</i> | 3 | 3 | 4 | 1 | 2 | 5 | 2 | 0 | 2 | 0 | 22 | 0,48% |
| <i>Patagioenas maculosa</i> | 1 | 0 | 2 | 0 | 1 | 0 | 2 | 0 | 0 | 0 | 6 | 0,13% |
| <i>Patagioenas picazuro</i> | 9 | 2 | 8 | 0 | 5 | 4 | 240 | 224 | 57 | 11 | 560 | 12,24% |
| <i>Zenaida auriculata</i> | 12 | 32 | 29 | 46 | 386 | 258 | 18 | 150 | 22 | 4 | 957 | 20,92% |
| <i>Myiopsitta monachus</i> | 15 | 12 | 12 | 8 | 23 | 9 | 14 | 28 | 25 | 31 | 177 | 3,87% |
| <i>Crotophaga ani</i> | 6 | 0 | 9 | 0 | 7 | 0 | 0 | 0 | 0 | 0 | 22 | 0,48% |
| <i>Guira guira</i> | 9 | 2 | 7 | 4 | 6 | 7 | 0 | 6 | 0 | 0 | 41 | 0,90% |
| <i>Piaya cayana</i> | 1 | 1 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 6 | 0,13% |

| | | | | | | | | | | | | |
|------------------------------------|---|---|---|---|---|---|---|---|----|----|----|-------|
| <i>Tyto alba</i> | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Athene cunicularia</i> | 2 | 0 | 2 | 1 | 0 | 0 | 2 | 1 | 0 | 0 | 8 | 0,17% |
| <i>Megascops choliba</i> | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 2 | 0,04% |
| <i>Bubo virginianus</i> | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 2 | 0,04% |
| <i>Hydropsalis torquata</i> | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0,04% |
| <i>Chlorostilbon lucidus</i> | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Hylocharis chrysura</i> | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 3 | 5 | 0,11% |
| <i>Megaceryle torquata</i> | 0 | 0 | 0 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 3 | 0,07% |
| <i>Chloroceryle amazona</i> | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 1 | 3 | 0,07% |
| <i>Chloroceryle americana</i> | 0 | 0 | 0 | 0 | 3 | 1 | 0 | 0 | 1 | 1 | 6 | 0,13% |
| <i>Colaptes campestris</i> | 4 | 2 | 3 | 0 | 5 | 4 | 2 | 3 | 5 | 2 | 30 | 0,66% |
| <i>Colaptes melanochloros</i> | 1 | 0 | 0 | 1 | 1 | 0 | 0 | 2 | 4 | 0 | 9 | 0,20% |
| <i>Melanerpes candidus</i> | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 5 | 0 | 0 | 7 | 0,15% |
| <i>Picumnus nebulosus</i> | 0 | 0 | 0 | 1 | 1 | 0 | 1 | 0 | 2 | 1 | 6 | 0,13% |
| <i>Veniliornis spilogaster</i> | 1 | 2 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 7 | 0,15% |
| <i>Thamnophilus caeruleus</i> | 0 | 1 | 2 | 1 | 2 | 2 | 4 | 0 | 1 | 2 | 15 | 0,33% |
| <i>Thamnophilus ruficapillus</i> | 5 | 2 | 2 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 10 | 0,22% |
| <i>Scytalopus pachecoi</i> | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Sittasomus griseicapillus</i> | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 2 | 0,04% |
| <i>Anumbius annumbi</i> | 2 | 0 | 6 | 0 | 0 | 0 | 1 | 3 | 2 | 8 | 22 | 0,48% |
| <i>Certhiaxis cinnamomeus</i> | 0 | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 4 | 0,09% |
| <i>Cranioleuca pyrrhophia</i> | 1 | 0 | 3 | 3 | 3 | 1 | 0 | 0 | 0 | 4 | 15 | 0,33% |
| <i>Lochmias nematura</i> | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 1 | 0 | 3 | 0,07% |
| <i>Phacellodomus striaticollis</i> | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 1 | 0 | 2 | 5 | 0,11% |
| <i>Limnortyx rectirostris</i> | 0 | 0 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 0,09% |
| <i>Furnarius rufus</i> | 6 | 4 | 6 | 5 | 4 | 6 | 9 | 5 | 22 | 13 | 80 | 1,75% |
| <i>Syndactyla rufosuperciliata</i> | 0 | 1 | 2 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 5 | 0,11% |
| <i>Synallaxis frontalis</i> | 0 | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 4 | 0,09% |
| <i>Synallaxis spixi</i> | 2 | 2 | 2 | 2 | 3 | 1 | 0 | 0 | 0 | 0 | 12 | 0,26% |
| <i>Phylloscartes ventralis</i> | 2 | 2 | 2 | 0 | 3 | 2 | 2 | 0 | 2 | 4 | 19 | 0,42% |
| <i>Poecilatriccus plumbeiceps</i> | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 2 | 0,04% |

| | | | | | | | | | | | | |
|----------------------------------|---|---|---|---|---|---|---|---|---|---|----|-------|
| <i>Elaenia obscura</i> | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0,04% |
| <i>Camptostoma obsoletum</i> | 2 | 1 | 2 | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 8 | 0,17% |
| <i>Serpophaga subcristata</i> | 5 | 2 | 2 | 0 | 2 | 3 | 0 | 2 | 2 | 4 | 22 | 0,48% |
| <i>Serpophaga nigricans</i> | 0 | 0 | 3 | 0 | 2 | 0 | 0 | 0 | 1 | 0 | 6 | 0,13% |
| <i>Xolmis cinereus</i> | 0 | 0 | 1 | 0 | 0 | 1 | 1 | 1 | 0 | 0 | 4 | 0,09% |
| <i>Xolmis dominicanus</i> | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Xolmis irupero</i> | 0 | 0 | 1 | 2 | 1 | 1 | 1 | 1 | 0 | 0 | 7 | 0,15% |
| <i>Hymenops perspicillatus</i> | 0 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 4 | 0,09% |
| <i>Knipolegus cyanirostris</i> | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 5 | 0,11% |
| <i>Pitangus sulphuratus</i> | 4 | 2 | 5 | 3 | 6 | 2 | 2 | 7 | 3 | 3 | 37 | 0,81% |
| <i>Machetornis rixosa</i> | 0 | 0 | 0 | 4 | 0 | 0 | 2 | 8 | 0 | 0 | 14 | 0,31% |
| <i>Lathrotriccus euleri</i> | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 2 | 0 | 4 | 0,09% |
| <i>Pyrocephalus rubinus</i> | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 7 | 0 | 7 | 0,15% |
| <i>Myiophobus fasciatus</i> | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Lessonia rufa</i> | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 0 | 0 | 0 | 4 | 0,09% |
| <i>Suiriri suiriri</i> | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 2 | 0,04% |
| <i>Empidonomus varius</i> | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0,02% |
| <i>Tyrannus melancholicus</i> | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 0 | 6 | 0 | 9 | 0,20% |
| <i>Myiarchus swainsoni</i> | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 0 | 2 | 0 | 7 | 0,15% |
| <i>Progne tapera</i> | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 0 | 0 | 0 | 5 | 0,11% |
| <i>Pygochelidon cyanoleuca</i> | 3 | 0 | 2 | 7 | 2 | 4 | 0 | 0 | 2 | 0 | 20 | 0,44% |
| <i>Stelgidopteryx ruficollis</i> | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0,04% |
| <i>Tachycineta leucorrhoa</i> | 0 | 0 | 4 | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 7 | 0,15% |
| <i>Cyclarhis gujanensis</i> | 1 | 5 | 1 | 1 | 1 | 2 | 0 | 0 | 0 | 0 | 11 | 0,24% |
| <i>Troglodytes musculus</i> | 4 | 2 | 3 | 1 | 8 | 2 | 1 | 0 | 3 | 3 | 27 | 0,59% |
| <i>Polioptila dumicola</i> | 2 | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 0 | 3 | 8 | 0,17% |
| <i>Turdus albicollis</i> | 0 | 2 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 3 | 0,07% |
| <i>Turdus amaurochalinus</i> | 4 | 9 | 5 | 2 | 7 | 5 | 3 | 0 | 1 | 0 | 36 | 0,79% |
| <i>Turdus rufiventris</i> | 6 | 8 | 8 | 3 | 9 | 3 | 1 | 2 | 2 | 2 | 44 | 0,96% |
| <i>Mimus saturninus</i> | 2 | 5 | 2 | 0 | 1 | 0 | 2 | 2 | 2 | 0 | 16 | 0,35% |
| <i>Anthus lutescens</i> | 0 | 0 | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 3 | 0,07% |

| | | | | | | | | | | | | |
|------------------------------------|----|----|----|----|----|-----|----|---|---|---|-----|-------|
| <i>Anthus hellmayri</i> | 1 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 3 | 0,07% |
| <i>Tangara sayaca</i> | 2 | 0 | 3 | 0 | 4 | 2 | 0 | 0 | 0 | 1 | 12 | 0,26% |
| <i>Tangara preciosa</i> | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 2 | 0,04% |
| <i>Stephanophorus diadematus</i> | 4 | 2 | 6 | 0 | 6 | 0 | 0 | 0 | 0 | 1 | 19 | 0,42% |
| <i>Pipraeidea bonariensis</i> | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 1 | 2 | 7 | 0,15% |
| <i>Saltator similis</i> | 0 | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 4 | 0,09% |
| <i>Saltator aurantiirostris</i> | 0 | 0 | 4 | 0 | 0 | 0 | 0 | 1 | 4 | 4 | 13 | 0,28% |
| <i>Paroaria coronata</i> | 4 | 2 | 2 | 2 | 2 | 2 | 6 | 6 | 8 | 4 | 38 | 0,83% |
| <i>Lanio cucullatus</i> | 2 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 0,07% |
| <i>Piranga flava</i> | 0 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 3 | 0 | 7 | 0,15% |
| <i>Zonotrichia capensis</i> | 12 | 14 | 13 | 16 | 27 | 8 | 7 | 2 | 5 | 6 | 110 | 2,40% |
| <i>Ammodramus humeralis</i> | 2 | 0 | 4 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 8 | 0,17% |
| <i>Sicalis flaveola</i> | 7 | 0 | 5 | 2 | 11 | 4 | 4 | 2 | 0 | 0 | 35 | 0,77% |
| <i>Sicalis luteola</i> | 0 | 0 | 13 | 0 | 9 | 0 | 0 | 0 | 0 | 0 | 22 | 0,48% |
| <i>Embernagra platensis</i> | 4 | 0 | 2 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 10 | 0,22% |
| <i>Poospiza cabanisi</i> | 6 | 3 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 0 | 15 | 0,33% |
| <i>Poospiza nigrorufa</i> | 3 | 0 | 2 | 0 | 2 | 0 | 4 | 0 | 0 | 2 | 13 | 0,28% |
| <i>Parula pitiayumi</i> | 3 | 4 | 2 | 1 | 6 | 2 | 2 | 1 | 0 | 1 | 22 | 0,48% |
| <i>Basileuterus culicivorus</i> | 2 | 2 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 6 | 0,13% |
| <i>Basileuterus leucoblepharus</i> | 4 | 4 | 2 | 2 | 6 | 2 | 0 | 0 | 5 | 2 | 27 | 0,59% |
| <i>Geothlypis aequinoctialis</i> | 2 | 0 | 3 | 0 | 3 | 1 | 0 | 0 | 0 | 0 | 9 | 0,20% |
| <i>Agelaioides badius</i> | 8 | 9 | 0 | 0 | 0 | 0 | 18 | 8 | 5 | 5 | 53 | 1,16% |
| <i>Cacicus chrysopterus</i> | 1 | 3 | 1 | 0 | 2 | 1 | 1 | 0 | 0 | 0 | 9 | 0,20% |
| <i>Gnorimopsar chopi</i> | 6 | 0 | 0 | 0 | 4 | 0 | 0 | 0 | 0 | 0 | 10 | 0,22% |
| <i>Molothrus bonariensis</i> | 10 | 2 | 37 | 9 | 54 | 290 | 16 | 7 | 0 | 5 | 430 | 9,40% |
| <i>Pseudoleistes guirahuro</i> | 0 | 0 | 8 | 0 | 12 | 0 | 0 | 0 | 0 | 0 | 20 | 0,44% |
| <i>Pseudoleistes virescens</i> | 0 | 1 | 0 | 0 | 0 | 0 | 40 | 0 | 0 | 0 | 41 | 0,90% |
| <i>Chrysomus ruficapillus</i> | 0 | 0 | 41 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 41 | 0,90% |
| <i>Xanthopsar flavus</i> | 14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 14 | 0,31% |
| <i>Icterus pyrrhopterus</i> | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 1 | 0 | 4 | 0,09% |
| <i>Sturnella superciliaris</i> | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0,02% |

Sporagra magellanica

9

2

4

0

18

7

3

0

0

1

44
4574

0,96%
100,00%